

DE QUERVAIN'S TENOSYNOVITIS

What is de Quervain's tenosynovitis?

De Quervain's tenosynovitis is a painful condition affecting the tendons located on the thumb side of your wrist. A tendon is a strong band of tissue that attaches muscle to bone. A sheath, or covering, surrounds the tendons that go to your thumb. Tenosynovitis is an irritation of this sheath.

How does it occur?

De Quervain's tenosynovitis usually occurs from overusing your thumb or wrist, especially in activities that move your thumb directly away from your wrist such as skiing or hammering.

What are the symptoms?

Symptoms may include:

- pain when you move your thumb or wrist
- pain when you make a fist
- swelling and tenderness on the thumb side of your wrist
- feeling or hearing creaking as the tendon slides through its sheath

How is it diagnosed?

Your healthcare provider will examine your wrist and thumb and find the areas that are tender and painful to move. An X-ray may be taken to be sure you don't have a broken bone.

How is it treated?

The initial treatment for de Quervain's tenosynovitis is a splint that will cover your wrist and thumb. It is important that you protect your thumb and wrist from activities that worsen your pain.

Treatment may also include:

- placing an ice pack on your thumb and wrist for 20 to 30 minutes every 3 or 4 hours until the pain goes away
- doing ice massage for 5 to 10 minutes several times a day
- taking an anti-inflammatory medicine such as ibuprofen (adults aged 65 years and older should not take non-steroidal anti-inflammatory medicine for more than 7 days without their healthcare provider's approval)
- having an injection of a medicine like cortisone

You will be given rehabilitation exercises to help speed your recovery and prevent the problem from returning.

When can I return to my sport or activity?


The goal of rehabilitation is to return you to your sport or activity as soon as is safely possible. If you return too soon you may worsen your injury, which could lead to permanent damage. Everyone recovers from injury at a different rate. Return to your activity will be determined by how soon your wrist recovers, not by how many days or weeks it has been since your injury occurred. In general, the longer you have symptoms before you start treatment, the longer it will take to get better.

You may return to your sport or activity when it is no longer painful to move your thumb or wrist. You may need to do activities wearing a supportive splint until you no longer have symptoms.

How can I prevent de Quervain's tenosynovitis?

Avoiding activities that overuse your thumb or wrist may prevent de Quervain's tenosynovitis.


DE QUERVAIN'S TENOSYNOVITIS




DE QUERVAIN'S TENOSYNOVITIS REHABILITATION EXERCISES

You may do all of these exercises when the initial pain is gone.

1. OPPOSITION STRETCH: Rest your hand on a table, palm up. Touch the tip of your thumb to the tip of your little finger. Hold this position for 6 seconds. Repeat 10 times.




OPPOSITION STRETCH



2. WRIST STRETCH: With one hand, help to bend the opposite wrist down by pressing the back of your hand and holding it down for 15 to 30 seconds. Next, stretch the hand back by pressing the fingers in a backward direction and holding it for 15 to 30 seconds. Keep your elbow straight during this exercise. Do 3 sets on each hand.


WRIST STRETCH

3. WRIST FLEXION: Hold a can or hammer handle in your hand with your palm facing up. Bend your wrist upward. Slowly lower the weight and return to the starting position. Do 3 sets of 10. Gradually increase the weight of the can or weight you are holding.



WRIST FLEXION

4. WRIST RADIAL DEVIATION STRENGTHENING: Put your wrist in the sideways position with your thumb up. Hold a can of soup or a hammer handle and gently bend your wrist up, with the thumb reaching toward the ceiling. Slowly lower to the starting position. Do not move your forearm throughout this exercise. Do 3 sets of 10.



WRIST RADIAL DEVIATION STRENGTHENING

5. WRIST EXTENSION: Hold a soup can or hammer handle in your hand with your palm facing down. Slowly bend your wrist upward. Slowly lower the weight down into the starting position. Do 3 sets of 10. Gradually increase the weight of the object you are holding.




WRIST EXTENSION

6. GRIP STRENGTHENING: Squeeze a rubber ball and hold for 5 seconds. Do 3 sets of 10.



GRIP STRENGTHENING



7. FINGER SPRING: Place a large rubber band around the outside of your thumb and the rest of your fingers. Open your fingers to stretch the rubber band. Do 3 sets of 10.

FINGER SPRING